arXiv:1302.5345v1 [math.NT] 21 Feb 2013

SOME IDENTITIES OF HIGHER-ORDER BERNOULLI, EULER AND HERMITE POLYNOMIALS ARISING FROM UMBRAL CALCULUS

DAE SAN KIM[1], TAEKYUN KIM[2], DMITRY V. DOLGY [3], AND SEOG-HOON RIM[4]

Abstract. In this paper, we study umbral calculus to have alternative ways of obtaining our results. That is, we derive some interesting identities of the higher-order Bernoulli, Euler and Hermite polynomials arising from umbral calculus to have alternative ways.

1. Introduction

As is well known, the Hermite polynomials are defined by the generating function to be

$$e^{2xt-t^2}=e^{H(x)t}=\sum_{n=0}^{\infty}H_n(x)\frac{t^n}{n!},\tag{1.1}$$

with the usual convention about replacing $H^n(x)$ by $H_n(x)$ (see [8, 10]). In the special case, $x=0$, $H_n(0)=H_n$ are called the *n-th Hermite numbers*. The Bernoulli polynomials of order r are given by the generating function to be

$$\left(\frac{t}{e^t-1}\right)^r e^{xt}=\sum_{n=0}^{\infty}B_n^{(r)}(x)\frac{t^n}{n!},\ (r\in\mathbb{R}).\tag{1.2}$$

From (1.2), the n-th Bernoulli numbers of order r are defined by $B_n^{(r)}(0)=B_n^{(r)}$ (see [1-16]). The higher-order Euler polynomials are also defined by the generating function to be

$$\left(\frac{2}{e^t+1}\right)^r e^{xt}=\sum_{n=0}^{\infty}E_n^{(r)}(x)\frac{t^n}{n!},\ (r\in\mathbb{R}),\tag{1.3}$$

and $E_n^{(r)}(0)=E_n^{(r)}$ are called the *n-th Euler numbers* of order r (see [1-16]).

The first Stirling number is given by

$$(x)_n=x(x-1)\cdots(x-n+1)=\sum_{l=0}^{n}S_1(n,k)x^l,\ \text{(see [6,13])},\tag{1.4}$$

and the second Stirling number is defined by the generating function to be

$$(e^t-1)^n=n!\sum_{l=n}^{\infty}S_2(l,n)\frac{t^l}{l!},\ \text{(see [6,9,13])}.\tag{1.5}$$

1991 *Mathematics Subject Classification.* 05A10, 05A19.

Key words and phrases. Bernoulli polynomial, Euler polynomial, Abel polynomial.

For $\lambda(\neq 1) \in \mathbb{C}$, the *Frobenius-Euler polynomials* are given by

$$\left(\frac{1-\lambda}{e^t-\lambda}\right)^r e^{xt} = \sum_{n=0}^{\infty} H_n^{(r)}(x|\lambda)\frac{t^n}{n!}, \ (r \in \mathbb{R}) \text{ (see [1,5]).} \tag{1.6}$$

In the special case, $x = 0$, $H_n^{(r)}(0|\lambda) = H_n^{(r)}(\lambda)$ are called the *n-th Frobenius-Euler numbers* of order r.

Let $\mathcal{F}$ be the set of all formal power series in the variable t over $\mathbb{C}$ with

$$\mathcal{F} = \left\{ f(t) = \sum_{k=0}^{\infty} \frac{a_k}{k!} t^k \;\middle|\; a_k \in \mathbb{C} \right\}.$$

Let us assume that $\mathbb{P}$ is the algebra of polynomials in the variable x over $\mathbb{C}$ and $\mathbb{P}^*$ is the vector space of all linear functionals on $\mathbb{P}$. $\langle L \mid p(x)\rangle$ denotes the action of the linear functional L on a polynomial $p(x)$ and we remind that the vector space structure on $\mathbb{P}^*$ is defined by

$$\begin{aligned} \langle L+M|p(x)\rangle &= \langle L|p(x)\rangle + \langle M|p(x)\rangle, \\ \langle cL|p(x)\rangle &= c\langle L|p(x)\rangle, \end{aligned}$$

where c is a complex constant (see [6, 9, 13]).

The formal power series $f(t) = \sum_{k=0}^{\infty} \frac{a_k}{k!} t^k \in \mathcal{F}$ defines a linear functional on $\mathbb{P}$ by setting

$$\langle f(t)|x^n\rangle = a_n, \text{ for all } n \geq 0 \text{ (see [6,9,13]).} \tag{1.7}$$

Then, by (1.7), we get

$$\left\langle t^k|x^n\right\rangle = n!\delta_{n,k}, \ (n, k \geq 0), \tag{1.8}$$

where $\delta_{n,k}$ is the Kronecker symbol (see [6, 9, 13]).

Let $f_L(t) = \sum_{k=0}^{\infty} \frac{\langle L|x^k\rangle}{k!} t^k$ (see [13]). For $f_L(t) = \sum_{k=0}^{\infty} \frac{\langle L|x^k\rangle}{k!} t^k$, we have $\langle f_L(t)|x^n\rangle = \langle L|x^n\rangle$. The map $L \mapsto f_L(t)$ is a vector space isomorphism from $\mathbb{P}^*$ onto $\mathcal{F}$. Henceforth, $\mathcal{F}$ will be thought of as both a formal power series and a linear functional. We shall call $\mathcal{F}$ the *umbral algebra*. The umbral calculus is the study of umbral algebra (see [6, 9, 13]).

The order $o(f(t))$ of the non-zero power series $f(t)$ is the smallest integer k for which the coefficient of t^k does not vanish. A series $f(t)$ having $o(f(t)) = 1$ is called a *delta series* and a series $f(t)$ having $o(f(t)) = 0$ is called an *invertible series*. Let $f(t)$ be a delta series and $g(t)$ be an invertible series. Then there exists a unique sequence $S_n(x)$ of polynomials such that $\left\langle g(t)f(t)^k|S_n(x)\right\rangle = n!\delta_{n,k}$ where $n, k \geq 0$. The sequence $S_n(x)$ is called *Sheffer sequence* for $(g(t), f(t))$, which is denoted by $S_n(x) \sim (g(t), f(t))$. By (1.7) and (1.8), we see that $\langle e^{yt}|p(x)\rangle = p(y)$. For $f(t) \in \mathcal{F}$ and $p(x) \in \mathbb{P}$, we have

$$f(t) = \sum_{k=0}^{\infty} \frac{\left\langle f(t)|x^k\right\rangle}{k!} t^k, \ p(x) = \sum_{k=0}^{\infty} \frac{\left\langle t^k|p(x)\right\rangle}{k!} x^k, \tag{1.9}$$

and, by (1.9), get

$$p^{(k)}(0) = \left\langle t^k|p(x)\right\rangle, \ \left\langle 1|p^{(k)}(x)\right\rangle = p^{(k)}(0). \tag{1.10}$$

Thus, from (1.10), we have

$$t^k p(x) = p^{(k)}(x) = \frac{d^k p(x)}{dx^k}. \tag{1.11}$$

In [6, 9, 13], we note that $\langle f(t)g(t)|p(x)\rangle = \langle g(t)|f(t)p(x)\rangle$.

For $S_n(x) \sim (g(t), f(t))$, we have

$$\frac{1}{g(\bar{f}(t))}e^{y\bar{f}(t)} = \sum_{k=0}^{\infty}\frac{S_k(y)}{k!}t^k, \text{ for all } y \in \mathbb{C}, \tag{1.12}$$

where $\bar{f}(t)$ is the compositional inverse of $f(t)$. For $S_n(x) \sim (g(t), f(t))$ and $r_n(x) = (h(t), l(t))$, let us assume that

$$S_n(x) = \sum_{k=0}^{n} C_{n,k} r_k(x), \text{ (see [6,9,13])}. \tag{1.13}$$

Then we have

$$C_{n,k} = \frac{1}{k!}\left\langle \frac{h(\bar{f}(t))}{g(\bar{f}(t))} l(\bar{f}(t))^k \middle| x^n \right\rangle, \text{ (see [13])}. \tag{1.14}$$

The equation (1.13) and (1.14) are called the alternative ways of Sheffer sequences.

In this paper, we study umbral calculus to have alternative ways of obtaining our results. That is, we derive some interesting identities of the higher-order Bernoulli, Euler and Hermite polynomials arising from umbral calculus to have alternative ways.

2. Some identities of higher-order Bernoulli, Euler and Hermite polynomials

In this section, we use umbral calculus to have alternative ways of obtaining our results. Let us consider the following Sheffer sequences:

$$E_n^{(r)}(x) \sim \left(\left(\frac{e^t+1}{2}\right)^r, t\right), H_n(x) \sim \left(e^{\frac{1}{4}t^2}, \frac{t}{2}\right). \tag{2.1}$$

Then, by (1.13), we assume that

$$E_n^{(r)}(x) = \sum_{k=0}^{n} C_{n,k} H_k(x). \tag{2.2}$$

From (1.14) and (2.2), we have

$$\begin{aligned}
C_{n,k} &= \frac{1}{k!}\left\langle \frac{e^{\frac{1}{4}t^2}}{\left(\frac{e^t+1}{2}\right)^r}\left(\frac{t}{2}\right)^k \middle| x^n \right\rangle \\
&= \frac{1}{k!2^k}\left\langle \left(\frac{2}{e^t+1}\right)^r e^{\frac{1}{4}t^2} \middle| t^k x^n \right\rangle \\
&= 2^{-k}\binom{n}{k}\left\langle \left(\frac{2}{e^t+1}\right)^r \middle| e^{\frac{1}{4}t^2} x^{n-k} \right\rangle \\
&= 2^{-k}\binom{n}{k}\left\langle \left(\frac{2}{e^t+1}\right)^r \middle| \sum_{l=0}^{\left[\frac{n-k}{2}\right]} \frac{1}{4^l l!} t^{2l} x^{n-k} \right\rangle \\
&= 2^{-k}\binom{n}{k}\sum_{l=0}^{\left[\frac{n-k}{2}\right]} \frac{1}{2^{2l}l!}(n-k)_{2l}\left\langle 1 \middle| \left(\frac{2}{e^t+1}\right)^r x^{n-k-2l} \right\rangle \\
&= 2^{-k}\binom{n}{k}\sum_{l=0}^{\left[\frac{n-k}{2}\right]} \frac{1}{2^{2l}l!}(n-k)_{2l} E_{n-k-2l}^{(r)} \\
&= n! \sum_{0\le l\le n-k,\ l:\text{even}} \frac{E_{n-k-l}^{(r)}}{\left(\frac{l}{2}\right)! 2^{k+l} k! (n-k-l)!}.
\end{aligned} \tag{2.3}$$

Therefore, by (2.2) and (2.3), we obtain the following theorem.

Theorem 2.1. *For $n \geq 0$, we have*

$$E_n^{(r)}(x) = n!\sum_{k=0}^{n}\left\{\sum_{0\le l\le n-k,\ l:even} \frac{E_{n-k-l}^{(r)}}{k!(n-k-l)!2^{k+l}\left(\frac{l}{2}\right)!}\right\} H_k(x).$$

Let us consider the following two Sheffer sequences:

$$B_n^{(r)}(x) \sim \left(\left(\frac{e^t-1}{t}\right)^r, t\right),\ H_n(x) \sim \left(e^{\frac{1}{4}t^2}, \frac{t}{2}\right). \tag{2.4}$$

Let us assume that

$$B_n^{(r)}(x) = \sum_{k=0}^{n} C_{n,k} H_k(x). \tag{2.5}$$

By (1.14) and (2.4), we get

$$\begin{aligned}
C_{n,k} &= \frac{1}{k!}\left\langle \frac{e^{\frac{1}{4}t^2}}{\left(\frac{e^t-1}{t}\right)^r}\left(\frac{t}{2}\right)^k \middle| x^n \right\rangle \\
&= 2^{-k}\binom{n}{k}\left\langle \left(\frac{t}{e^t-1}\right)^r \middle| \sum_{l=0}^{\infty}\left(\frac{1}{4}\right)^l \frac{1}{l!}t^{2l}x^{n-k} \right\rangle \\
&= 2^{-k}\binom{n}{k}\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{1}{l!4^l}(n-k)_{2l}\left\langle \left(\frac{t}{e^t-1}\right)^r \middle| x^{n-k-2l} \right\rangle \\
&= 2^{-k}\binom{n}{k}\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{(n-k)!}{l!2^{2l}(n-k-2l)!}\left\langle 1 \middle| \left(\frac{t}{e^t-1}\right)^r x^{n-k-2l} \right\rangle \\
&= 2^{-k}\binom{n}{k}\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{(n-k)!}{l!2^{2l}(n-k-2l)!}B_{n-k-2l}^{(r)} \\
&= n!\sum_{0\le l\le n-k,\ l:\text{even}}\frac{B_{n-k-l}^{(r)}}{k!(n-k-l)!2^{k+l}\left(\frac{l}{2}\right)!}.
\end{aligned} \tag{2.6}$$

Therefore, by (2.5) and (2.6), we obtain the following theorem.

Theorem 2.2. *For $n \ge 0$, we have*

$$B_n^{(r)}(x) = n!\sum_{k=0}^{n}\left\{\sum_{0\le l\le n-k,\ l:even}\frac{B_{n-k-l}^{(r)}}{k!(n-k-l)!2^{k+l}\left(\frac{l}{2}\right)!}\right\}H_k(x).$$

Consider

$$H_n^{(r)}(x|\lambda) \sim \left(\left(\frac{e^t-\lambda}{1-\lambda}\right)^r, t\right), H_n(x) \sim \left(e^{\frac{1}{4}t^2}, \frac{t}{2}\right). \tag{2.7}$$

Let us assume that

$$H_n^{(r)}(x|\lambda) = \sum_{k=0}^{n} C_{n,k}H_k(x). \tag{2.8}$$

By (1.14), we get

$$\begin{aligned}
C_{n,k} &= \frac{1}{k!}\left\langle \frac{e^{\frac{1}{4}t^2}}{\left(\frac{e^t-\lambda}{1-\lambda}\right)^r}\left(\frac{t}{2}\right)^k \middle| x^n \right\rangle \\
&= \frac{1}{k!2^k}\left\langle \left(\frac{1-\lambda}{e^t-\lambda}\right)^r e^{\frac{1}{4}t^2} \middle| t^k x^n \right\rangle \\
&= 2^{-k}\binom{n}{k}\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{(n-k)_{2l}}{l!4^l}\left\langle 1 \middle| \left(\frac{1-\lambda}{e^t-\lambda}\right)^r x^{n-k-2l}\right\rangle \\
&= n!\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{H^{(r)}_{n-k-2l}(\lambda)}{l!2^{2l+k}(n-k-2l)!k!} \\
&= n!\sum_{0\le l\le n-k,\ l:\text{even}}\frac{H^{(r)}_{n-k-l}(\lambda)}{\left(\frac{l}{2}\right)!2^{k+l}(n-k-l)!k!}.
\end{aligned} \tag{2.9}$$

Therefore, by (2.8) and (2.9), we obtain the following theorem.

Theorem 2.3. *For $n \geq 0$, we have*

$$H^{(r)}_n(x|\lambda) = n!\sum_{k=0}^{n}\left\{\sum_{0\le l\le n-k,\ l:even}\frac{H^{(r)}_{n-k-l}(\lambda)}{k!(n-k-l)!\left(\frac{l}{2}\right)!2^{k+l}}\right\}H_k(x).$$

Let us assume that

$$H_n(x) = \sum_{k=0}^{n} C_{n,k}E^{(r)}_k(x). \tag{2.10}$$

From (1.14), (2.1) and (2.10), we have

$$\begin{aligned}
C_{n,k} &= \frac{1}{k!}\left\langle \frac{\left(\frac{e^{2t}+1}{2}\right)^r}{e^{\frac{1}{4}(2t)^2}}(2t)^k \middle| x^n \right\rangle \\
&= \frac{1}{k!}\left\langle \frac{\left(\frac{e^{t}+1}{2}\right)^r}{e^{\frac{1}{4}t^2}}t^k \middle| (2x)^n \right\rangle \\
&= \frac{1}{k!}2^n\left\langle \left(\frac{e^t+1}{2}\right)^r e^{-\frac{1}{4}t^2} \middle| t^k x^n \right\rangle \\
&= 2^n\binom{n}{k}\left\langle \left(\frac{e^t+1}{2}\right)^r \middle| \sum_{l=0}^{\infty}\frac{(-1)^l}{l!4^l}t^{2l}x^{n-k}\right\rangle \\
&= 2^{n-r}\binom{n}{k}\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{(-1)^l}{l!2^{2l}}(n-k)_{2l}\left\langle \left(e^t+1\right)^r \middle| x^{n-k-2l}\right\rangle \\
&= \frac{1}{2^r}\sum_{j=0}^{r}\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{\binom{n}{k}\binom{r}{j}2^k(-1)^l(n-k)!}{l!(n-k-2l)!}(2j)^{n-k-2l}.
\end{aligned} \tag{2.11}$$

Therefore, (2.10) and (2.11), we obtain the following theorem.

Theorem 2.4. *For $n \geq 0$, we have*

$$H_n(x) = \frac{1}{2^r}\sum_{k=0}^{n}\left\{\sum_{j=0}^{r}\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{\binom{n}{k}\binom{r}{j}2^k(-1)^l(n-k)!(2j)^{n-k-2l}}{l!(n-k-2l)!}\right\}E_k^{(r)}(x).$$

Note that $H_n(x) \sim \left(e^{\frac{1}{4}t^2}, \frac{t}{2}\right)$. Thus, we have

$$e^{\frac{1}{4}t^2}H_n(x) \sim \left(1, \frac{t}{2}\right), \text{ and } (2x)^n \sim \left(1, \frac{t}{2}\right). \tag{2.12}$$

From (2.12), we have

$$e^{\frac{1}{4}t^2}H_n(x) = (2x)^n \Leftrightarrow H_n(x) = e^{-\frac{1}{4}t^2}(2x)^n. \tag{2.13}$$

By (2.11) and (2.13), we also see that

$$\begin{aligned}
C_{n,k} &= \frac{1}{k!}\left\langle \left.\frac{\left(\frac{e^{2t}+1}{2}\right)^r}{e^{\frac{1}{4}(2t)^2}}(2t)^k \right| x^n\right\rangle \\
&= \frac{1}{k!}\left\langle \left.\left(\frac{e^t+1}{2}\right)^r t^k \right| e^{-\frac{1}{4}t^2}(2x)^n\right\rangle \\
&= \frac{1}{k!2^r}\left\langle (e^t+1)^r \middle| t^k H_n(x)\right\rangle = \frac{1}{2^r}\binom{n}{k}2^k\sum_{j=0}^{r}\binom{r}{j}H_{n-k}(j).
\end{aligned} \tag{2.14}$$

Therefore, by (2.10) and (2.14), we obtain the following theorem.

Theorem 2.5. *For $n \geq 0$, we have*

$$H_n(x) = \frac{1}{2^r}\sum_{k=0}^{n}\binom{n}{k}2^k\left[\sum_{j=0}^{r}\binom{r}{j}H_{n-k}(j)\right]E_k^{(r)}(x).$$

Let us assume that

$$H_n(x) = \sum_{k=0}^{n}C_{n,k}B_k^{(r)}(x). \tag{2.15}$$

From (1.14), (2.4) and (2.15), we have

$$\begin{aligned}
C_{n,k} &= \frac{1}{k!}\left\langle \left.\frac{\left(\frac{e^{2t}-1}{2t}\right)^r}{e^{\frac{1}{4}(2t)^2}}(2t)^k \right| x^n\right\rangle \\
&= \frac{1}{k!}\left\langle \left.\frac{\left(\frac{e^t-1}{t}\right)^r}{e^{\frac{1}{4}t^2}}t^k \right| (2x)^n\right\rangle \\
&= \frac{1}{k!}\left\langle \left.\left(\frac{e^t-1}{t}\right)^r t^k \right| e^{-\frac{1}{4}t^2}(2x)^n\right\rangle.
\end{aligned} \tag{2.16}$$

From (2.13) and (2.16), we have

$$C_{n,k} = \frac{1}{k!}\left\langle \left.\left(\frac{e^t-1}{t}\right)^r t^k \right| H_n(x)\right\rangle. \tag{2.17}$$

For $r > n$, by (1.5) and (2.17), we get

$$\begin{aligned} C_{n,k} &= \frac{1}{k!}\left\langle (e^t-1)^k \left| \left(\frac{e^t-1}{t}\right)^{r-k} H_n(x) \right.\right\rangle \\ &= \frac{1}{k!}\left\langle (e^t-1)^k \left| \sum_{l=0}^{n} \frac{(r-k)!}{(l+r-k)!} S_2(l+r-k, r-k) t^l H_n(x) \right.\right\rangle \\ &= \frac{1}{k!}\sum_{l=0}^{n} \frac{(r-k)!}{(l+r-k)!} S_2(l+r-k,r-k) 2^l (n)_l \left\langle (e^t-1)^k | H_{n-l}(x)\right\rangle \\ &= \frac{1}{k!}\sum_{l=0}^{n} \frac{(r-k)!}{(l+r-k)!} S_2(l+r-k,r-k) 2^l \frac{n!}{(n-l)!} \sum_{j=0}^{k} \binom{k}{j}(-1)^{k-j} H_{n-l}(j) \\ &= n! \sum_{j=0}^{k}\sum_{l=0}^{n} \frac{(r-k)! S_2(l+r-k,r-k)(-1)^{k-j}\binom{k}{j} 2^l H_{n-l}(j)}{(l+r-k)!k!(n-l)!}. \end{aligned} \tag{2.18}$$

Therefore, by (2.15) and (2.18), we obtain the following theorem.

Theorem 2.6. *For $r > n \geq 0$,we have*

$$H_n(x) = n! \sum_{k=0}^{n} \left\{ \sum_{j=0}^{k}\sum_{l=0}^{n} \frac{(r-k)! S_2(l+r-k,r-k)(-1)^{k-j}\binom{k}{j} 2^l H_{n-l}(j)}{(l+r-k)!k!(n-l)!} \right\} B_k^{(r)}(x).$$

Let us assume that $r \leq n$. For $0 \leq k < r$, by (2.18), we get

$$C_{n,k} = n! \sum_{j=0}^{k}\sum_{l=0}^{n} \frac{(r-k)! S_2(l+r-k,r-k)(-1)^{k-j}\binom{k}{j} 2^l H_{n-l}(j)}{(l+r-k)!k!(n-l)!}. \tag{2.19}$$

For $r \leq k \leq n$, by (2.17), we get

$$\begin{aligned} C_{n,k} &= \frac{1}{k!}\sum_{j=0}^{r}\binom{r}{j}(-1)^{r-j}\left\langle e^{jt} | D^{k-r} H_n(x)\right\rangle \\ &= \frac{2^{k-r} n!}{k!(n-k+r)!}\sum_{j=0}^{r}\binom{r}{j}(-1)^{r-j} H_{n-k+r}(j). \end{aligned} \tag{2.20}$$

Therefore, by (2.15), (2.19) and (2.21), we obtain the following theorem.

Theorem 2.7. *For $n \geq r$, we have*

$$\begin{aligned} H_n(x) =& n! \sum_{k=0}^{r-1} \left\{ \sum_{j=0}^{k}\sum_{l=0}^{n} \frac{(r-k)! S_2(l+r-k,r-k)(-1)^{k-j}\binom{k}{j} 2^l H_{n-l}(j)}{(l+r-k)!k!(n-l)!} \right\} B_k^{(r)}(x) \\ &+ n! \sum_{k=r}^{n} \left\{ \sum_{j=0}^{r} \frac{(-1)^{r-j}\binom{r}{j} 2^{k-r} H_{n-k+r}(j)}{k!(n-k+r)!} \right\} B_k^{(r)}(x). \end{aligned}$$

Let us assume that

$$H_n(x) = \sum_{k=0}^{n} C_{n,k} H_k^{(r)}(x|\lambda). \tag{2.21}$$

Then, by (1.14), (2.7) and (2.21), we get

$$\begin{aligned} C_{n,k} &= \frac{1}{k!}\left\langle \frac{\left(\frac{e^{2t}-\lambda}{1-\lambda}\right)^r}{e^{\frac{1}{4}(2t)^2}}(2t)^k \middle| x^n \right\rangle \\ &= \frac{1}{k!}\left\langle \frac{\left(\frac{e^{t}-\lambda}{1-\lambda}\right)^r}{e^{\frac{1}{4}t^2}}t^k \middle| (2x)^n \right\rangle \\ &= \frac{1}{k!}\left\langle \left(\frac{e^{t}-\lambda}{1-\lambda}\right)^r t^k \middle| e^{-\frac{1}{4}t^2}(2x)^n \right\rangle. \end{aligned} \tag{2.22}$$

By (2.13) and (2.22), we get

$$\begin{aligned} C_{n,k} &= \frac{1}{k!}\left\langle \left(\frac{e^{t}-\lambda}{1-\lambda}\right)^r t^k \middle| H_n(x) \right\rangle = \frac{1}{k!(1-\lambda)^r}\left\langle (e^t-\lambda)^r \middle| t^k H_n(x)\right\rangle \\ &= \frac{\binom{n}{k}2^k}{(1-\lambda)^r}\sum_{j=0}^{r}\binom{r}{j}(-\lambda)^{r-j}\left\langle e^{jt} | H_{n-k}(x)\right\rangle \\ &= \frac{\binom{n}{k}2^k}{(1-\lambda)^r}\sum_{j=0}^{r}\binom{r}{j}(-\lambda)^{r-j}H_{n-k}(j). \end{aligned} \tag{2.23}$$

Therefore, by (2.21) and (2.23), we obtain the following theorem.

Theorem 2.8. *For $n \geq 0$, we have*

$$H_n(x) = \frac{1}{(1-\lambda)^r}\sum_{k=0}^{n}\binom{n}{k}2^k\left[\sum_{j=0}^{r}\binom{r}{j}(-\lambda)^{r-j}H_{n-k}(j)\right]H_k^{(r)}(x|\lambda).$$

Remark. From (2.22), we have

$$\begin{aligned} C_{n,k} &= \frac{1}{k!}\left\langle \frac{\left(\frac{e^{t}-\lambda}{1-\lambda}\right)^r}{e^{\frac{1}{4}t^2}}t^k \middle| (2x)^n \right\rangle = \frac{2^n}{k!}\left\langle \left(\frac{e^{t}-\lambda}{1-\lambda}\right)^r e^{-\frac{1}{4}t^2} \middle| t^k x^n \right\rangle \\ &= \frac{(n)_k}{k!}2^n\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{(-1)^l}{l!4^l}\left\langle \left(\frac{e^{t}-\lambda}{1-\lambda}\right)^r \middle| t^{2l}x^{n-k}\right\rangle \\ &= \frac{\binom{n}{k}2^n}{(1-\lambda)^r}\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{(-1)^l}{l!2^{2l}}(n-k)_{2l}\left\langle \left(e^t-\lambda\right)^r \middle| x^{n-k-2l}\right\rangle \\ &= \frac{1}{(1-\lambda)^r}\sum_{j=0}^{r}\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{\binom{n}{k}\binom{r}{j}2^k(-1)^l(-\lambda)^{r-j}(n-k)!(2j)^{n-k-2l}}{l!(n-k-2l)!}. \end{aligned} \tag{2.24}$$

Thus, by (2.21) and (2.24), we get

$$H_n(x) = \frac{1}{(1-\lambda)^r}\sum_{k=0}^{n}\left\{\sum_{j=0}^{r}\sum_{l=0}^{\left[\frac{n-k}{2}\right]}\frac{\binom{n}{k}\binom{r}{j}2^k(-1)^l(-\lambda)^{r-j}(n-k)!(2j)^{n-k-2l}}{l!(n-k-2l)!}\right\}H_k^{(r)}(x|\lambda).$$

References

[1] S. Araci, M. Acikgoz, *A note on the Frobenius-Euler numbers and polynomials associated with Bernstein polynomials*, Adv. Stud. Contemp. Math. **22** (2012), no. 3, 399-406. .

[2] S. Araci, D. Erdal, J. Seo, *A study on the fermionic p-adic q-integral representation on* $\mathbb{Z}_p$ *associated with weighted q-Bernstein and q-Genocchi polynomials*, Abstr. Appl. Anal. **2011** (2011), Art. ID 649248, 10 pp.

[3] A. Bayad,*Modular properties of elliptic Bernoulli and Euler functions*, Adv. Stud. Contemp. Math. **20** (2010), no. 3, 389-401.

[4] M. Can, M. Cenkci, V. Kurt and Y. Simsek, *Twisted Dedekind type sums associated with Barnes' type multiple Frobenius-Eulerl-functions*, Adv. Stud. Contemp. Math. **18** (2009), no. 2, 135-160.

[5] L. Carlitz, *The product of two Eulerian polynomials*, Math. Mag. **368** (1963) 37-41.

[6] R. Dere and Y. Simsek, *Applications of umbral algebra to some special polynomials*, Adv. Stud. Contemp. Math. **22** (2012) 433-438.

[7] D. Ding and J. Yang, *Some identities related to the Apostol-Euler and Apostol-Bernoulli polynomials*, Adv. Stud. Contemp. Math. **20** (2010), no. 1, 7-21.

[8] D. S. Kim, T. Kim, S.-H. Rim and S.-H. Lee, *Hermite polynomials and their applications associated with Bernoulli and Euler Numbers*, Discrete Dynamics in Nature and Society, (2012), Article ID 974632, 13 pages doi:10.1155/2012/974632

[9] D. S. Kim and T. Kim, *Some identities of Frobenius-Euler polynomials arising from umbral calculus*, Adv. Difference Equ. 2012, 2012:196 doi:10.1186/1687-1847-2012-196.

[10] T. Kim, J.Choi, Y.H. Kim and C. S. Ryoo, *On q-Bernstein and q-Hermite polynomials*, Proc. Jangjeon Math. Soc. **14**(2011), no. 2, 215-221.

[11] H. Ozden, I. N. Cangul and Y. Simsek, *Remarks on q-Bernoulli numbers associated with Daehee numbers*, Adv. Stud. Contemp. Math. **18** (2009), no. 1, 41-48.

[12] S.-H. Rim, J. Joung, J.-H. Jin and S.-J. Lee, *A note on the weighted Carlitzs type q-Euler numbers and q-Bernstein polynomials*, Proc. Jangjeon Math. Soc. **15** (2012) 195-201.

[13] S. Roman, *The umbral calculus*, Dover Publ. Inc. New York, 2005.

[14] C. Ryoo, *Some relations between twisted q-Euler numbers and Bernstein polynomials*, Adv. Stud. Contemp. Math. **21** (2011), no. 2, 217-223.

[15] Y. Simsek, *Special functions related to Dedekind-type DC-sums and their applications*, Russ. J. Phys. **17** (2010) 495-508.

[16] Y. Simsek, *Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions*, Adv. Stud. Contemp. Math. **16** (2008), no. 2, 251-278.

[1] DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, REPUBLIC OF KOREA.
E-mail address: `dskim@sogang.ac.kr`

[2] DEPARTMENT OF MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA.
E-mail address: `tkkim@kw.ac.kr`

[3] HANRIMWON, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA.
E-mail address: `dgekw2011@gmail.com`

[4] DEPARTMENT OF MATHEMATICS EDUCATION, KYUNGPOOK NATIONAL UNIVERSITY, TAEGU 702-701, REPUBLIC OF KOREA.
E-mail address: `shrim@knu.ac.kr`